

- Northeast Germany – Landscape Hydrology – Vegetation – Virtual water

**Gunnar Lischeid and Marco Natkhnin**

## The Potential of Land-Use Change to Mitigate Water Scarcity in Northeast Germany – a Review

*Das Potential von Landnutzungsänderungen zur Minderung der  
Wasserknappheit in Nordostdeutschland – ein Überblick*

With 6 Figures and 1 Table

Climate change is expected to increase water scarcity in northeast Germany. Land-use change is one of the options of mitigation that is intensely discussed in this region. This review aims at giving a compilation of existing data and modelling studies in order to investigate the potential and the limits of the land-use change approach.

### 1. Introduction

Water is essential for life on earth as well as for human civilisation. It is a basic food for man, a primary means of agricultural and silvicultural production and an important means of industrial production. Rivers and channels form waterways for transportation of goods. Freshwater bodies and wetlands play a major role for biodiversity as well as for nutrient and contaminant retention in the landscape. Last but not least, lakes and streams form scenic landscapes that are highly attractive for recreation and tourism, especially in northeast Germany. Thus, any issues that are related to landscape hydrology attract wide interest in the public.

Lakes, wetlands and streams cover a substantial fraction of the lowlands in northeast Germany, that is, in the federal states of Mecklenburg-Vorpommern, Brandenburg, Berlin, and in the north of Saxony-Anhalt. Compared to the rest of Germany, annual rainfall is very low and air temperature during the growing season rather high, resulting in high evapotranspiration and small groundwater recharge. During the last three decades, substantially decreasing lake water and groundwater levels have been observed in many parts of the region (*Gerstengarbe et al. 2003*). As a consequence, lakes have been shrinking and small lakes, streams and wetlands have dried out. This has caused concern in the public as well as in water resources management, environmental

agencies, and politics. Various options have been brought forward and discussed for mitigating water scarcity in northeast Germany.

Major consumption of incoming rainfall is by transpiration from agricultural fields and forests. However, substantial differences have been found between different types of land use as well as between different agricultural and silvicultural management strategies. Thus many mitigation strategies focus on reducing water loss via transpiration by land-use optimisation. This topic was addressed, among others, in the NEWAL-NET (*Natkhan 2011*) project which was established in 2005. It aimed at developing a management strategy for a sustainable development of forests in the lowlands of northeast Germany with regard to the predicted climate change, i.e., above all, decreasing availability of soil water and groundwater during the growing season. Some of the management options outlined there are based on different water use by different species. This paper benefits substantially from the work undertaken in this project.

Besides, it provides a review on a variety of experimental and modeling studies, scenario analyses and theoretical frameworks that have been developed both for northeast Germany and for other parts of the world. The latter applies especially to the virtual water concept that has been developed for irrigation cropping systems in Asia but is now discussed to be applied to northeast Germany as well (*Lischeid 2010*). The objective of the paper is to assess the opportunities and limitations of mitigating water scarcity by land-use and land management optimisation in northeast Germany.


## **2. Study Region: Geology, Soils, Climate, Hydrology, Agriculture**

The German lowlands are part of the North German-Polish Basin which has been subject to subsidence and recurrent marine transgressions

since the end of the Palaeozoic. Consequently, marine sediments like clay, sand and lime mud have been deposited. Total thickness of these sediments, especially in the northern part of the region, is a few hundred metres (*Lippstreu et al. 1997, Stackebrandt and Manhenke 2002*). However, the landform configuration has been shaped more recently during the Pleistocene with repeated advances of glaciers from the northeast and subsequent retreats. The terminal moraines now form the hilly regions between the extended lowlands of the former ice marginal valleys. The latters stretch from the southeast to the northwest, or from east to west, respectively, along the former flow direction of the melting water that discharged to the North Sea (*Fig. 1*). Altitude of these lowlands usually is a few tens of metres above sea level, whereas it exceeds 100 m above sea level in parts of the terminal moraines region. Consequently, topographic gradients and river water flow velocities are very low in the lowlands. Due to the young age of the landscape and the very shallow topographic gradients at larger scales, drainage basins are not well developed. Thus, many lakes and wetlands have been connected to the stream network only by man-made ditches.

Soils are mostly sandy in most parts of the region. More loamy soils are encountered in fluvial sediments and in the vicinity of ground moraines, especially in the northern part of the region. In the lowlands peaty soils and fens are abundant. However, both soils and deeper sediments are characterised by very high spatial heterogeneities and short spatial correlation lengths.

Compared to other parts of Germany, precipitation is fairly low and air temperature rather high. Annual mean rainfall in Brandenburg and Berlin is about 610 mm per year (*Gerstengarbe et al. 2003, Lahmer and Pfützner 2003, Dannowski and Steidl 2000*). Annual mean temperature 1951-2000 was between 7.8°C and 9.5°C (*Gerstengarbe et al. 2003*), with lower temperatures in the north com-




*Fig. 1 Topography of Brandenburg (by courtesy of Uwe Heinrich)  
Topographie Brandenburgs (zu Verfügung gestellt von Uwe Heinrich)*

pared to the south. Evapotranspiration amounts to 510 mm per year roughly, leaving 100 mm per year as runoff. About 80 mm per year occurs as groundwater recharge (Gerstengarbe et al. 2003, Lahmer and Pfützner 2003, Dannowski and Steidl 2000). However, groundwater recharge is negative in the lowlands where the groundwater level is close to the surface. In contrast, it amounts to

more than 200 mm per year in parts of the hilly regions (Lahmer and Pfützner 2003).

Especially on sandy soils with very small water holding capacity, evapotranspiration highly depends on depth to groundwater. It is less than 2 m in 39 % of the area in Brandenburg and Berlin, and more than 5 m in 60 % of the area (Lahmer



*Fig. 2 Land use in the federal states of Mecklenburg-Vorpommern and Brandenburg in 2004. Data: Deggau 2006 / Landnutzung in Mecklenburg-Vorpommern und Brandenburg 2004. Daten: Deggau 2006*

and Pfützner 2003). This bimodal distribution reflects the fraction of lowlands in the former ice margin valleys on the one hand, and the hilly regions on the other hand, where even tree roots do not have access to groundwater resources.

Most of northeast Germany is covered by the two federal states of Mecklenburg-Vorpommern and Brandenburg. Here, more than half of the area is used by agriculture (Fig. 2), where arable land prevails. Another 30 % is covered by forests. Surface water covers another 4.4 %. Settlements, transport installations and other infrastructure make up another 11 % of the area – and nearly 70 % in Berlin (Deggau 2006).

### 3. The Virtual Water Concept

Water consumption by plants competes with alternative water usage, e.g., as drinking water, as means of industrial production, or as the basic resource of wetland and freshwater habitats. The water content of agricultural products is usually

only a minute fraction of the total water consumption during the growth of the plants. The production of 1 kg of crops requires some thousands of litres of water. Thus, any calculation of water consumption for meat production has to account for that of fodder crops, resulting in even higher figures (Hoekstra 2003, Rockström 2003). Total water consumption during the production process, including water content of the products, is summarized as the “virtual water” content of the products. It is calculated by dividing total water consumption by crop yield (Hoekstra 2003).

This concept was developed by Allan (1993, 1994) to account for water consumption in extensively irrigated fields in the Middle East. Production and export of, e.g. cotton, thus implies export of virtual water which should be considered in economic assessments of resources allocation in arid regions (Allan 1994, Liu et al. 2009).

The virtual water concept has gained great attention and has been considerably extended since it was first formulated, e.g. it has turned out to be

useful to differentiate between different types of water resources: “Green water” means rainwater that infiltrates into the soil and feeds the plants. In contrast, “blue water” denotes irrigation water that needs to be pumped from groundwater or surface water. Third, “grey water” is waste water that people want to get rid of, which, however, can be used for irrigation (Liu et al. 2009).

This distinction addresses a crucial point, that is, the opportunity costs of the water used (Liu et al. 2009): Opportunity costs usually are the highest for blue water, because this may be used for other purposes, e.g., for drinking water supply. For grey water, in contrast, opportunity costs may even be negative, when no alternative usage is possible without additional expensive treatment. In addition, using grey water for irrigation makes efficient use of the water’s nutrient load and of the purification potential of the soil.

On the other hand, however, increasing irrigation may even lower the virtual water content: Applying irrigation to agricultural crops at the end of the ripening season can ensure ripening of the crops that otherwise die off. Thus, a rather small increase of water consumption via irrigation may result in an overproportional increase of crop yield (Rockström 2003, Liu et al. 2007).

#### **4. The North East Germany Case Study: Land Use and Land Management Options**

In northeast Germany, by far the largest fraction of rainwater is consumed by plant transpiration in forests, arable land and grassland. About 510 mm per year out of 610 mm per year of precipitation is lost by evapotranspiration, where transpiration dominates by far (Gerstengarbe et al. 2003, Lahmer and Pfützner 2003, Dannowski and Steidl 2000). At a global scale, the largest fraction of water resources is used for agricultural production (Rockström 2003). Different crops and species differ substantially with respect to rain consump-

tion. Thus, land-use change and land management are considered as a major tool for reducing water consumption in water-scarce regions.

However, it has to be taken into account that other factors play an important role for evapotranspiration as well. For example, effects of geomorphology and scale effects must be considered (Cuevas et al. 2006, Strayer et al. 2003, Herlihy et al. 1998). Besides, the water holding capacity of the soil, surface runoff depending on soil properties as well as on soil tillage and rainfall intensity, or seasonal patterns of rainfall account for much of the heterogeneities in the available data. In addition, nonlinear interactions have to be accounted for. For example, water holding capacity and rooting depth are negligible as long as precipitation or irrigation occur evenly throughout the growing season. In contrast, soils may develop a hydrophobic topsoil layer that impedes rainwater infiltration, depending on the properties of soil organic material and the length of dry periods (Doerr et al. 2000). Comprehensive data sets and sophisticated models are required to account for these effects.

##### *4.1 Lakes and wetlands*

Northeast Germany is characterised by a large number of lakes, kettle holes (Kaletka 1996) and wetlands. Evapotranspiration from these water bodies usually is highly overproportional. For example, Herbst and Kappen (1993) found that evapotranspiration of a reed stand in a north German lake was about 187 % compared to evaporation from the lake surface. Behrendt et al. (2001) observed an exponential increase of annual evapotranspiration with decreasing depth to groundwater at Paulinenaue, northwest of Berlin. Here, annual evapotranspiration of a reed stand (*Phragmites australis*) was about 2000 mm per year, which was more than threefold the mean annual precipitation. Similarly, evapotranspiration of wetland plants determined *in situ* by Dannowski and Balla (2004) and Müller et al.

Tab. 1 Runoff for various types of land use in Brandenburg and Berlin under present-day climatic conditions (Dannowski and Steidl 2000) / Abfluss für verschiedene Landnutzungstypen in Brandenburg und Berlin unter aktuellen klimatischen Bedingungen (Dannowski and Steidl 2000)

	Entire year	Dormant season	Growing season
	[mm yr <sup>-1</sup> ]		
<b>Uncultivated land</b>	309	96	213
<b>Urban areas</b>	244	94	150
<b>Agricultural land</b>	127	75	52
<b>Forest</b>	56	94	-38
<b>Surface water</b>	-103	133	-236

(2005) exceeded 1000 mm per year, which was roughly equal to twice the annual precipitation.

#### 4.2 Forests

In general, evapotranspiration is highest in coniferous forests, less in deciduous forests, and even less on arable land and grassland. Differences are mainly due to varying length of the period of physiological activity, varying aerodynamic roughness of the vegetation structure, and varying rooting depth (Brown et al. 2005). Schindler et al. (2008) investigated different land-use forms at Müncheberg, east of Berlin, from 1995 through 2005. Annual seepage flux under arable land and grassland was nearly identical with 175 mm per year and 174 mm per year, respectively. In contrast, mean annual seepage flux in a pine forest (*Pinus sylvestris*) was only 15 mm per year, that is, less than 10 % compared to the aforementioned.

Correspondingly, the effect of land use on the water balance is reflected in numerous modeling studies. Converting forests into arable and grassland increased runoff in the adjacent stream by 9 % in the model study by Fohrer et al. (2001). Converting greenland to forest and arable land decreased runoff by 3 %.

Runoff was modelled for the federal states of Brandenburg and Berlin by Dannowski and Steidl (2000), using the ABIMO model. The difference between uncultivated land on the one hand and surface water on the other hand was about 400 mm per year (Tab. 1). In this study, evapotranspiration exceeded precipitation during the growing season in forests and lakes. For the latter, the surplus of precipitation during the dormant season could not compensate for high water losses in summer. However, runoff exhibited substantial spatial variability, depending on local meteorological conditions. Thus, e.g., even some of the pine forests exhibited negative water budgets where precipitation was fairly low.

Past land-use changes are likely to have exerted a major impact on water budgets as well. Based on a comprehensive data set, Bork et al. (1998) conclude that the proportion of forested area in Germany decreased from 90 % to 15 % between the early Middle Ages and the beginning of the 14<sup>th</sup> century. The corresponding increase of runoff was assessed to be 120 mm per year. Given that annual runoff is only about 100 mm per year today in northeast Germany, this change must have had consid-




Fig. 3 Simulated annual mean groundwater recharge for different land-use scenarios and soils in the Schorfheide region 1952-2007 (source: Natkhan 2011) / Simulierte jährliche Grundwasserneubildung für verschiedene Landnutzungsszenarien und Böden in der Schorfheide 1952-2007 (Quelle: Natkhan 2011)


erable effects on lake water levels, extension of wetland areas and river runoff.

Natkhan (2011) studied the effect of land-use change in the Schorfheide region for different soils, using the WaSiM-ETH model. Meteorological data from the region covering the 1952-2007 period were used, and typical soils were considered. Mean annual groundwater recharge amounted to 45 mm per year for pine forests, 114 mm per year for beech stands, and 215 mm per year for grassland (Fig. 3). Differences between types of land use were most pronounced for coarse sand overlying loam, illustrating the interplay between the water holding capacity of the soil and the rooting depth.

Correspondingly, Bolte et al. (2002) investigated the effect of replacing the actual forests in Germany by natural forests with a higher proportion of deciduous trees. In their scenario, the result-

ing increase of seepage flux was only 16 mm per year for all of Germany. However, the effect was much larger in northeast Germany due to limited water resources. In another model study, Wattenbach et al. (2007) found that replacing all pine forests in the federal state of Brandenburg by deciduous forests would increase mean groundwater recharge by 4.5 %. Mey and Pfützner (2008) conclude from their results that replacing pine stands by a beech forest in the vicinity of the Luch lake, 30 km southeast of Berlin, would increase the groundwater level by 0.4 m.

Fröhlich et al. (2008) investigated the effect of various measures on the groundwater level in a 442 km<sup>2</sup> area in the Schorfheide region in Brandenburg, using a complex hydrological model. Reducing pine forests by 8 % and increasing oak forests by 10 %, as suggested in forest management plans, hardly had any impact on groundwater levels. In contrast, replacing the existing forests by potential



*Fig. 4* Water budgets of different forests at a site close to Eberswalde with 620 mm per year mean annual precipitation. “P” denotes pine, “B” beech. In addition, tree age [years] is given in brackets. Data provided by Müller 2002 / Wasserbilanz für verschiedene Forstbestände nahe Eberswalde mit 620 mm/a Jahresniederschlag. „P“ steht für Kiefer, „B“ für Buche. Außerdem ist das Baumalter [Jahre] in Klammern angegeben. Datengrundlage: Müller 2002

natural vegetation with a high proportion of oak and beech increased mean groundwater recharge by 63 mm per year. To that end, pine forests had to be reduced by 58 %, beech forests to be increased by 34 %, and mixed pine and beech forests by 21 %.


In addition to tree species, tree age and understory vegetation play an important role. A comprehensive experimental data set was presented by Müller (2002). In general, an increase of evapotranspiration with tree age was observed for pine and beech stands. However, evapotranspiration decreased in a later stage in the pine stand – a process which had not been observed in the beech stand (*Fig. 4*). In addition, the mixing ratio of pine and beech had nonlinear effects on the total stand evapotranspiration.

The extensive deforestation during and after World War II, and intensive afforestation thereafter, is reflected by the age distribution of pine

forests in Brandenburg. In addition, the fraction of trees less than 20 years old is rather small due to enhanced planting of deciduous trees (Müller 2007). This would result in a long-term increase of groundwater recharge, if boundary conditions remained stable. On the other hand, understory vegetation changed considerably since the 1970s. Due to increased atmospheric deposition, a nitrophilic understory established in many pine forests (Hofmann 1995), presumably increasing both evapotranspiration and interception.

#### 4.3 Crops

Corresponding to different forest types, crops differ substantially with respect to water use, too. This information can be used to develop and optimise adaptation strategies in the water-scarce region of northeast Germany. However, differences between crops are usually smaller




*Fig. 5 Assessment of annual water use by different crops in Germany in two studies / Abschätzung des jährlichen Wasserverbrauchs durch unterschiedliche Feldfrüchte in Deutschland in zwei verschiedenen Studien*

compared to those between forests on the one hand, and arable land or grassland on the other hand. As a consequence, additional effects of soil type, depth to groundwater level, soil tillage, seasonal patterns of precipitation, fertilisation and pest control need to be considered as well (*Liu et al. 2007, Hoekstra and Hung 2002*).

Due to limited precipitation and sandy soils with poor water holding capacity water is one of the key minimum factors for plant growth and plant production in northeast Germany. Investigating a typical crop vegetation scheme including winter wheat, sugar beet, summer barley and pea in the Uckermark region in the study period 1992-1999, *Schindler et al. (2001)* found that the total amount of summer

precipitation (April-September) explained most of the interannual variance of crop production. In contrast, fertilisation had hardly any effect on crop yield. However, this is not necessarily valid for other regions of Germany.

These side effects are rarely considered when crop water use is assessed, e.g., within the framework of the virtual water concept. Correspondingly, the data given in the literature exhibit substantial variance. For example, figures for annual water use by wheat and barley in Germany differed by up to 150 mm per year in two different studies (*Fig. 5*). In general, differences between crops in each of these studies are smaller compared to those between these studies. In another assessment by *Hoekstra (2003)* water use by maize varies by a



*Fig. 6* Water use divided by crop yield by different cereals on non-irrigated fields (mean and range) compiled by Rockström (2003) for the tropics (rice; corn, i.e. maize) or the mid-latitudes (other cereals), respectively. Data provided by (1) Rockström et al. (1999) and Doorenbos and Kassam (1986)  
*Auf den Ertrag bezogener Wasserverbrauch verschiedener Getreidearten ohne künstliche Bewässerung (jeweils Mittelwert und Spannweite) für die Tropen (Reis, Mais) und mittleren Breiten (alle anderen Getreidearten), zusammengestellt von Rockström (2003) auf Basis der Daten von (1) Rockström et al. (1999) und Doorenbos und Kassam (1986)*

factor up to four. However, those figures apply to different regions with different climate.

Differences between single crops are partly due to varying length of the required growing period. Thus, water use by vegetables and lettuce is very low, and that of sugar beet very high (*Fig. 5*). On the other hand, different metabolic pathways in so-called C3 and C4 plants play a role. Only the latter, like, e.g., maize, can assimilate carbon during the night and during dry periods when stomata are closed. In contrast, C3 plants have to take up CO<sub>2</sub> during daytime when transpiration water loss is high.

Besides these effects, there is some doubt whether the maximum possible water use efficiency,

related to crop yield, does even differ systematically between crops and regions. In the data set compiled by Rockström (2003) water use efficiency for various cereals differed by a factor of five (*Fig. 6*). However, he did not find any systematic differences between climatic regions. He concludes that higher transpiration in dry and warm regions is compensated for by higher productivity. The author ascribes lower water use efficiency found by numerous studies in dry warm regions to higher non-productive water loss in gappy crops or yield losses, e.g., due to water scarcity at the end of the growing period. Considering only optimal growing conditions, water use efficiency seems to converge to 1300–1500 l kg<sup>-1</sup> for various cereals irrespective of the climate region (Rockström 2003). Rockström concludes that this figure seems

to reflect a global physiological minimum, resulting in an absolute minimum water use for food production of  $1300 \text{ m}^3$  per year per capita.

Assessing the global water budget based on a crop growth model, *Liu et al.* (2009) conclude that about 81 % of global crop production is based on green water, that is, rainwater. This figure is close to the 85 % given by *Rost et al.* (2008). In fact, irrigation is restricted to about 20 % of arable land (*Rockström* 2003). Here, more efficient techniques like drip irrigation or rainwater harvesting should be applied which would reduce the total amount of blue water considerably (*Liu et al.* 2009). Evaporation from the soil can be further reduced by, e.g., mulch-till and hedges that reduce wind velocity.

Virtual water import through the import of food and fodder to Germany exceeds virtual water export by far. According to *Hoekstra* and *Hung* (2002) and *Chapagain* and *Hoekstra* (2003) net import of virtual water amounts to  $12.2 \cdot 10^9$  or  $13.1 \cdot 10^9 \text{ m}^3$  per year, respectively. In contrast, net export is only  $1 \cdot 10^9 \text{ m}^3$  per year according to *Zimmer* and *Renault* (2003) which means that import is about twelve or thirteen times the value of export.

*Sonnenberg et al.* (2009) assess the total annual consumption of virtual water by agricultural products in Germany, including meat, to be equal to  $117.6 \cdot 10^6 \text{ m}^3$  per year. Roughly half of this virtual water is imported, with coffee, cocoa, oil seeds, cotton, pork, soybeans and beef being the most important products. Agricultural products sum up to 73.3 % of the total virtual water import in Germany. The rest is ascribed to industrial products and drinking water withdrawal (*Sonnenberg et al.* 2009).

## 5. Climate Change Effects

Water resources management has to take into account the expected climate change of the

next decades. Depending on the selected scenario of greenhouse gas emissions, global climate models differ with respect to the long-term increase of air temperature. Greenhouse gas emissions measured over the last decades correspond to the more pessimistic scenarios for which the models predict an increase of air temperature of about 1K in the next 50 years (*Solomon et al.* 2009).

Spatial resolution of the global circulation models is not sufficient to assess regional effects. Thus, results of the global models need to be scaled down by regional climate models. A dynamic scaling approach is followed by the REMO model (*Jacob et al.* 2008). It is based on a weather forecast model with a detailed simulation of physical processes in the atmosphere at a 10 km resolution, using the output of global models as boundary conditions. Alternative approaches assume that statistical relationships between large-scale patterns, as predicted by global circulation models, and small-scale patterns will remain stable in the next decades and can be used for down-scaling the results of the global circulation models. To that end, the models STAR (*Werner* and *Gerstengarbe* 1997) and WETTREG (*Spekat et al.* 2007) have been developed for Germany.

All the regional models predict a substantial increase of temperature over the next 50 to 100 years in northeast Germany based on the moderate and pessimistic scenarios of future greenhouse gas emissions. Trends are less clear for precipitation. However, most results indicate that precipitation will tend to decrease during the growing season and to increase during the dormant season. In general, the models agree in that evapotranspiration during the summer period is likely to increase whilst groundwater recharge and runoff will most likely decrease in the next decades (*Jacob et al.* 2008, *Spekat et al.* 2007, *Wechsung* 2005, *Gerstengarbe et al.* 2003, *Suckow et al.* 2002).

*Gerstengarbe* et al. (2003) assessed the effects of climate change on crop yield in northeast Germany. Neither an increase of irrigation nor any possible progress with regard to pest control or plant breeding was considered. Observed crop yield during the 1980-1990 period was compared with model results for the 2040-2050 period, based on a moderate greenhouse gas emission increase scenario. Temperature was predicted to increase by 1.7 K, and precipitation to decrease by 52 mm per year compared to 1980-1990. As a consequence, wheat yield was expected to decrease by 17 %, and maize yield to increase. Taking into account the effect of increasing CO<sub>2</sub> concentration, the predicted wheat yield decreased by 10 %, and maize and barley yield increased by 8 % and 7 %, respectively.

Similar results are reported by *Wechsung* et al. (2008). In their model, the increase of CO<sub>2</sub> partial pressure will compensate for water scarcity for the next 40 years in East Germany. However, the drought risk will increase, especially in Brandenburg. Additional effects of increasing frequencies of extreme events or pest infestation on crop yield might play a role but can only be speculated about. In contrast, fire risk is almost certainly expected to increase, especially in the pine forests in Brandenburg (*Gerstengarbe* and *Werner* 1997). This provides another strong argument for converting pine forests into mixed stands or deciduous forests.

## 6. Conclusions

Landscape hydrology in northeast Germany is susceptible to minor changes of the water budget, due to the large proportion of area covered by lakes and wetlands on the one hand, and low present-day groundwater recharge and runoff generation on the other hand. Due to climate change, water scarcity is expected to increase further within the next decades. Thus, water

resources management considers different land-use options in order to adapt to and to mitigate water scarcity in northeast Germany.

Wetlands and small lakes exhibit the highest rates of evapotranspiration, usually exceeding local precipitation. Thus, these systems provide the largest water sinks per area in northeast Germany. Among the major land-use types in northeast Germany, pine forests rank second with respect to annual water use. Converting them into deciduous forests is considered a major option with regard to increasing groundwater recharge and stream runoff. However, due to long rotation periods, forest conversion is a very slow process compared to the velocity of climate change. Whether this option may compensate for the future increase of water scarcity or even increase water budgets at the landscape scale depends to a large extent on the details of forest conversion and the local boundary conditions.

Groundwater recharge and runoff generation from grassland and arable land is substantially higher compared to that of forests. Various crops are generally assumed to differ in this respect. However, except for vegetables with a short growing period, there seems to be a physiological minimum water use of cereals in general, irrespective of crop species or climate. That maximum water use efficiency is warranted only given optimal soil tillage, fertilisation, pest control and irrigation, if necessary. Apart from the latter, which still does not pay off for most crops grown in the region today, it seems that crop production in northeast Germany is already fairly close to the water use optimum.

Transpiration is the price to pay for vegetation cover. Correspondingly, groundwater recharge is highest on uncultivated land. From a purely water resources management aspect, the most efficient way to increase groundwater recharge and stream runoff in northeast Germany would be to convert lakes and wetlands into fallows.

However, this would be at odds with numerous competing objectives and is not considered a serious water management option. Besides, wetlands and lakes cover only a minor fraction of the area in northeast Germany.

In contrast, most projects focus on converting pine forests into deciduous forests. In fact, this conversion will yield a substantial decrease of annual evapotranspiration. This option is generally preferred, also with regard to its impacts on forest fire and infestation risk, and the corresponding benefits with respect to biodiversity. Converting pine forests into arable land or grassland would reduce evapotranspiration even more. This is hardly enforceable given conservation issues, existing property rights and legal reasons. However, it is worth considering an alternative patchy mixture of low-intensity pasture and deciduous forests. This so-called “semipen pasture” system (*Reisinger* 2005) has been considered mainly with respect to biodiversity and nature conservation issues so far. Based on the available data about water use of pine forests, deciduous forests and grassland in the mid-latitudes, such a strategy deserves being considered as a vital water resources management option. However, non-linear effects in these heterogeneous systems need to be studied first. For example, due to the “oasis effect”, small forest stands within arable fields transpire much more per area during hot summer days compared to extended forest.

Alternative approaches are based on using water resources unused so far. Grey water, i.e. waste water, is being used for irrigation in many parts of the world. As a side effect, enhanced microbial activity and the associated decomposition of organic pollutants in the topsoil could at least partly be a surrogate for waste water treatment plants. In addition, nutrients contained in waste water could be used by the plants without any pre-treatment. Last but not least, a surplus of irrigated grey water might increase groundwater

recharge and stream runoff and increase the water level in lakes and wetlands further downstream. Thus, the opportunity costs of grey water irrigation would be negative. Due to legal restrictions that account for the risk of ground-water pollution, however, waste water irrigation is allowed only in very few cases in Germany. However, it should be considered to use waste water after initial treatment. Both groundwater levels and stream water quality would benefit from this and save further costly treatment in waste water plants. In fact, facing the predicted increase of water scarcity problems in northeast Germany, new concepts are required to adapt to and to mitigate water scarcity.

## 7. References

- Allan, J.A.* 1993: Fortunately there are Substitutes for Water otherwise our Hydro-Political Futures would be Impossible. – In: ODA (ed.): Priorities for Water Resources Allocation and Management. – London: 13-26
- Allan, J.A.* 1994: Overall Perspectives on Countries and Regions. – In: *Rogers, P. and P. Lydon* (eds.): Water in the Arab World: Perspectives and Prognoses. – Cambridge: 65-100
- Behrendt, A., G. Schalitz, L. Müller, G. Mundel und D. Hözel* 2001: Untersuchungen zur Niedermoorenrenaturierung. – In: 9. Gumpensteiner Lysimetertagung, 24. und 25. April 2001, Bundesanstalt für alpenländische Landwirtschaft Gumpenstein. – Irdning: 141-144
- Bolte, A., B. Wolff und S. Anders* 2002: Tiefensickerung in Abhängigkeit von realer und natürlicher Waldbedeckung: Eine bundesweite Modellstudie. – In: Funktionen des Waldes und Aufgaben der Forstwirtschaft in Verbindung mit dem Landschaftswasserhaushalt. Praxiskolloquium der Landesforststalt Eberswalde am 25. Oktober 2001 anlässlich des 150. Geburtstages des Boden- und Standortskundlers Prof. Dr. Emil Ramann. – Eberswalder Forstliche Schriftenreihe 15. – Potsdam: 77-86
- Bork, H.-R., H. Bork, C. Dalchow, B. Faust, H.-P. Piorr und T. Schatz* 1998: Landschaftsentwicklung in Mitteleuropa: Wirkungen des Menschen auf Landschaften. – Gotha, Stuttgart

- Brown, A.E., L. Zhang, T.A. McMahon, A.W. Western and R.A. Vertessy* 2005: A Review of Paired Catchment Studies for Determining Changes in Water Yield Resulting from Alterations in Vegetation. – *Journal of Hydrology* **310** (1-4): 28-61
- Chapagain, A.K. and A.Y. Hoekstra* 2003: Virtual Water Trade: A Quantification of Virtual Water Flows between Nations in Relation to International Trade of Livestock and Livestock Products. – In: *Hoekstra, A.Y.* (ed.): Virtual Water Trade. Proceedings of the International Expert Meeting on Virtual Water Trade. – Value of Water Research Report Series **12**. – Delft: 48-76
- Chapagain, A.K. and A.Y. Hoekstra* 2004: Water Footprints of Nations. Volume 2: Appendices. – Value of Water Research Report Series **16**. – Delft
- Cuevas, J.G., D. Soto, I. Arismende, M. Pino, A. Lara and C. Oyarzún* 2006: Relating Land Cover to Stream Properties in Southern Chilean Watersheds: Trade-off between Geographic Scale, Sample Size, and Explanatory Power. – *Biogeochemistry* **81** (3): 313-329
- Dannowski, R. und D. Balla* 2004: Wasserhaushalt und geohydrologische Situation einer vernässten Niedermoorfläche mit Schilfanzbau in Nordost-Brandenburg. – *Archiv für Naturschutz und Landschaftsforschung* **43** (2): 27-40
- Dannowski, R. und J. Steidl* 2000: Modellierung des Gebietswasserhaushaltes mit dem Modell ABIMO. – In: Landesumweltamt Brandenburg (Hrsg.): Flächen-deckende Modellierung von Wasserhaushaltsgrößen für das Land Brandenburg. – Studien und Tagungsberichte, Landesumweltamt Brandenburg **27**. – Potsdam: 24-40
- Deggau, M.* 2006: Nutzung der Bodenfläche. Flächen-erhebung 2004 nach Art der tatsächlichen Nutzung. – *Wirtschaft und Statistik* (3): 212-219
- Doerr, S., R.A. Shakesby and R.P.D. Walsh* 2000: Soil Water Repellency: Its Causes, Characteristics and Hydro-Geomorphological Significance. – *Earth-Science Reviews* **51**: 33-65
- Doorenbos, J. and A.H. Kassam* 1986: Yield Response to Water. – *FAO Irrigation and Drainage Paper* **33**. – Rome
- Fohrer, N., S. Haverkamp, K. Eckhardt and H.G. Frede* 2001: Hydrologic Response to Land Use Changes on the Catchment Scale. – *Physics and Chemistry of the Earth Part B* **26** (7-8): 577-582
- Fröhlich, K., B. Hasch und M. Dembeck* 2008: Machbarkeitsstudie „Stabilisierung des Landschaftswasserhaushaltes der Schorfheide“. – Berlin
- Gerstengarbe, F.-W. und P.C. Werner* 1997: Waldbrandentwicklung im Land Brandenburg. – *Der Wald* **7**: 392-394
- Gerstengarbe, F.-W., F. Badeck, F. Hattermann, V. Krysanova, W. Lahmer, P. Lasch, M. Stock, F. Suckow, F. Wechsung und P.C. Werner* 2003: Studie zur klimatischen Entwicklung im Land Brandenburg bis 2055 und deren Auswirkungen auf den Wasserhaushalt, die Forst- und Landwirtschaft sowie die Ableitung erster Perspektiven. – *PIK-Report* **83**. – Potsdam
- Herbst, M. und L. Kappen* 1993: Die Rolle des Schilfs im standörtlichen Wasserhaushalt eines norddeutschen Sees. – *Phytocoenologia* **23**: 51-64
- Herlihy, A.T., J.L. Stoddard and C.B. Johnson* 1998: The Relationships between Stream Chemistry and Watershed Land Cover Data in the Mid-Atlantic Region, U.S. – *Water, Air, and Soil Pollution* **105** (1-2): 377-386
- Hoekstra, A.Y.* 2003: Virtual Water: An Introduction. – In: *Hoekstra, A.Y.* (ed.): Virtual Water Trade. Proceedings of the International Expert Meeting on Virtual Water Trade. – Value of Water Research Report Series **12**. – Delft: 13-23
- Hoekstra, A.Y. and P.Q. Hung* 2002: A Quantification of Virtual Water Flows Between Nations in Relation to International Crop Trade. – Value of Water Research Report Series **11**. – Delft
- Hofmann, G.* 1995: Zur Wirkung von Stickstoffeinträgen auf die Vegetation nordostdeutscher Kiefernwaldungen. – In: IMA-Querschnittseminar Wirkungskomplex Stickstoff und Wald. – Umweltbundesamt, Texte **28**. – Berlin: 131-140
- Jacob, D., H. Göttel, S. Kotlarski, P. Lorenz und K. Sieck* 2008: Klimaauswirkungen und Anpassung in Deutschland. Phase 1: Erstellung regionaler Klimaszenarien für Deutschland. – Abschlussbericht zum UFOPLAN-Vorhaben 204 41 13. – Dessau
- Kaletka, T.* 1996: Die Problematik der Sölle (Klein-hohlformen) im Jungmoränengebiet Nordostdeutschlands. – *Naturschutz und Landespflege in Brandenburg*, Sonderheft Sölle: 4-12
- Lahmer, W. und B. Pfützner* 2003: Orts- und zeitdiscrete Ermittlung der Sickerwassermenge im Land Brandenburg auf der Basis flächendeckender Wasserhaushaltsberechnungen. – *PIK-Report* **85**. – Potsdam

- Lippstreu, L., N. Hermsdorf und A. Sonntag* 1997: Geologische Übersichtskarte des Landes Brandenburg 1:300.000. – Erläuterungen. – Potsdam
- Lischeid, G.* 2010: Landschaftswasserhaushalt in der Region Berlin-Brandenburg. – Berlin-Brandenburgische Akademie der Wissenschaften, Interdisziplinäre Arbeitsgruppe Globaler Wandel – Regionale Entwicklung, Diskussionspapier 01/2010. – Berlin
- Liu, J., H. Yang and A. Zehnder* 2007: Simulation of Crop Water Relations on Large Scales with High Spatial Resolutions. – In: *Van Bers, C., D. Petry and C. Pahl-Wostl* (eds.): Global Assessments: Bridging Scales and Linking to Policy. Report on the Joint TIAS-GWSP Workshop held at the University of Maryland University College, Adelphi, USA, 10 - 11 May 2007. – GWSP Issues in Global Water System Research 2. – Bonn: 44-48
- Liu, J., H. Yang and A. Zender* 2009: Global Consumptive Water Use for Crop Production: The Importance of Green Water and Virtual Water. – Water Resources Research 45: W05428, doi: 10.1029/2007WR006051
- Mey, S. und B. Pfützner* 2008: Analyse zur Minde rung von Wassermangelsituationen mit einem gekoppelten Oberflächen-Grundwassерmodell. – Poster, Tag der Hydrologie 2008. – Hannover. – Online available: <http://www.iww.uni-hannover.de/tdh2008/Poster/Mey.pdf>, 16/12/2009
- Müller, J.* 2002: Verdunstung der Baumvegetation und Tiefenversickerung in Kiefern-, Buchen- und Eichenökosystemen in Abhängigkeit von Boden- und Bestandesstruktur und von der Witterung. – In: *Anders, S., W. Beck, A. Bolte, G. Hofmann, M. Janssen, U.-K. Krakau und J. Müller*: Ökologie und Vegetation der Wälder Nordostdeutschlands. Einfluss von Niederschlagsarmut und erhöhtem Stickstoffeintrag auf Kiefern-, Eichen- und Buchen-Wald- und Forstökosysteme des nordostdeutschen Tieflandes. – Remagen-Oberwinter: 113-122
- Müller, K.* 2007: Die aktuelle Verbreitung der Kiefer in Brandenburg. – In: *Kätzel, R., K. Möller, S. Löfler, J. Engel und K. Liero* (Hrsg.): Die Kiefer im nordostdeutschen Tiefland – Ökologie und Bewirtschaftung. – Eberswalder Forstliche Schriftenreihe 32. – Eberswalde: 9-13
- Müller, L., A. Behrendt, G. Schalitz und U. Schindler* 2005: Above Ground Biomass and Water Use Efficiency of Crops at Shallow Water Tables in a Temperate Climate. – Agricultural Water Management 75 (2): 117-136
- Natkhin, M.* 2011: Modellgestützte Analyse der Einflüsse von Veränderungen der Waldwirtschaft und des Klimas auf den Wasserhaushalt grundwasser-abhängiger Landschaftselemente. – Dissertation, Universität Potsdam
- Reisinger, E.* 2005: Ausgewählte naturschutzfachliche und sozioökonomische Anforderungen für die Etablierung großflächiger Weidesysteme. – In: *Finck, P., W. Härdtle, B. Redecker und U. Riecker* (Hrsg.): Weidelandschaften und Wildnisgebiete – Vom Experiment zur Praxis. Referate und Ergebnisse der gleichnamigen Fachtagung an der Universität Lüneburg vom 23. bis zum 26. September 2003. – Schriftenreihe für Landschaftspflege und Naturschutz 78. – Münster: 469-489
- Rockström, J.* 2003: Water for Food and Nature in Drought-Prone Tropics: Vapour Shift in Rain-Fed Agriculture. – Philosophical Transactions of the Royal Society London, Biological Sciences 358: 1997-2009
- Rockström, J., L. Gordon, M. Falkenmark, C. Folke and M. Engvall* 1999: Linkages Among Water Vapor Flows, Food Production, and Terrestrial Ecosystem Services. – Conservation Ecology 3 (2): 1-28
- Rost, S., D. Gerten, A. Bondeau, W. Lucht, J. Rohwer und S. Schaphoff* 2008: Agricultural Green and Blue Water Consumption and its Influence on the Global Water System. – Water Resources Research 44 (9): 1-17
- Schindler, U., M. Wolff und G. Kühn* 2001: Lysimeterstudie zum Einfluss von Düngung und Bewirtschaftung auf die Ertragsbildung, den Wasserhaushalt und die Nährstoffauswaschung im Trockengebiet der Uckermark. – Journal of Plant Nutrition and Soil Science 164 (6): 697-703
- Schindler, U., L. Müller, F. Eulenstein and R. Dannowski* 2008: A Long-Term Hydrological Soil Study on the Effects of Soil and Land Use on Deep Seepage Dynamics in Northeast Germany. – Archives of Agronomy and Soil Science 54 (5): 451-463
- Solomon, S., D. Qin, M. Manning, R.B. Alley, T. Berntsen, N.L. Bindoff, Z. Chen, A. Chidthaisong, J.M. Gregory, G.C. Hegerl, M. Heimann, B. Hewitson, B.J. Hoskins, F. Joos, J. Jouzel, V. Kattsov, U. Lohmann, T. Matsuno, M. Molina, N. Nicholls, J. Overpeck, G. Raga, V. Ramaswamy, J. Ren, M. Rusticucci, R. Somerville, T.F. Stocker, P. Whet-*

- ton, R.A. Wood and D. Wratt 2007: Technical Summary. – In: Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.): Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. – Cambridge, New York*
- Sonnenberg, A., A. Chapagain, M. Geiger und D. August 2009: Der Wasser-Fußabdruck Deutschlands. Woher stammt das Wasser, das in unseren Lebensmitteln steckt? – Frankfurt/Main*
- Spekat, A., W. Enke and F. Kreienkamp 2007: Neuentwicklung von regional hoch aufgelösten Wetterlagen für Deutschland und Bereitstellung regionaler Klimaszenarios auf der Basis von globalen Klimasimulationen mit dem Regionalisierungsmodell WETTREG auf der Basis von globalen Klimasimulationen mit ECHAM5/MPI-OM T63L31 2010 bis 2100 für die SRES-Szenarios B1, A1B und A2. – Umweltbundesamt. – Dessau*
- Stackebrandt, W. und V. Manhenke 2002: Atlas zur Geologie von Brandenburg im Maßstab 1:1 000 000. – Landesamt für Geowissenschaften und Rohstoffe Brandenburg. – Kleinmachnow*
- Strayer, D.L., R.E. Beighley, L.C. Thompson, S. Brooks, C. Nilsson, G. Pinay and R.J. Naiman 2003: Effects of Land Cover on Stream Ecosystems: Roles of Empirical Models and Scaling Issues. – Ecosystems 6 (5): 407–423*
- Suckow, F., P. Lasch und F.-W. Badeck 2002: Auswirkungen von Klimaveränderungen auf die Grundwasserneubildung. – In: Funktionen des Waldes und Aufgaben der Forstwirtschaft in Verbindung mit dem Landschaftswasserhaushalt. Praxiskolloquium der Landesforstanstalt Eberswalde am 25. Oktober 2001 anlässlich des 150. Geburtstages des Boden- und Standortskundlers Prof. Dr. Emil Ramann. – Eberswalder Forstliche Schriftenreihe 15. – Potsdam: 36-44*
- Wattenbach, M., M. Zebisch, F. Hattermann, P. Gottschalk, H. Goemann, P. Kreins, F. Badeck, P. Lasch, F. Suckow and F. Wechsung 2007: Hydrological Impact Assessment of Afforestation and Change in Tree-Species Composition: A Regional Case Study for the Federal State of Brandenburg (Germany). – Journal of Hydrology 346 (1-2): 1-17*
- Wechsung, F. 2005: Herausforderungen des globalen Wandels für die Elbe-Region. – In: Wechsung, F., A. Becker und P. Gräfe (Hrsg.): Auswirkungen des globalen Wandels auf Wasser, Umwelt und Gesellschaft im Elbegebiet. – Konzepte für die nachhaltige Entwicklung einer Flusslandschaft 6. – Berlin: 3-57*
- Wechsung, F., A. Lüttger und F.F. Hattermann 2008: Projektionen zur klimabedingten Änderung der Erträge von einjährigen Sommer- und Winterkulturen des Ackerlandes am Beispiel von Silomais und Winterweizen. – In: Wechsung, F., F.-W. Gerstengarbe, P. Lasch und A. Lüttger (Hrsg.): Die Ertragsfähigkeit ostdeutscher Ackerflächen unter Klimawandel. – PIK-Report 112. – Potsdam: 18-32*
- Werner, P.C. and F.W. Gerstengarbe 1997: Proposal for the Development of Climate Scenarios. – Climate Research 8 (3): 171-182*
- Zimmer, D. and D. Renault 2003: Virtual Water in Food Production and Global Trade: Review of Methodological Issues and Preliminary Results. – In: Hoekstra, A.Y. (ed.): Virtual Water Trade. – Proceedings of the International Expert Meeting on Virtual Water Trade. – Value of Water Research Report Series 12. – Delft: 93-109*
- Summary: The Potential of Land-Use Change to Mitigate Water Scarcity in Northeast Germany – a Review**
- Water scarcity has become a major topic in northeast Germany. Annual groundwater recharge is already close to zero in parts of the region. Moreover, groundwater and lake water levels have decreased substantially during the last three decades and are likely to decrease further. By far the largest consumption of rainwater is due to transpiration by agricultural plants and forests. Thus, many mitigation strategies aim at maximising water use efficiency by land-use adaptation and agricultural or forestry management optimisation. This paper gives a review about the opportunities of these approaches, with both the actual and the predicted future climatic boundary conditions in northeast Germany in mind.

*Zusammenfassung: Das Potential von Landnutzungsänderungen zur Minderung der Wasserknappheit in Nordostdeutschland – ein Überblick*

Wasserknappheit wird in Nordostdeutschland zunehmend zu einem Problem. Die aktuelle jährliche Grundwasserneubildung ist in Teilen der Region schon nahe Null. Zusätzlich weisen Grund- und Seewasserspiegel in den letzten 30 Jahren verbreitet deutliche Rückgänge auf und werden voraussichtlich in Zukunft weiter absinken. Der mit Abstand größte Teil des Wasserverbrauchs in der Landschaft entfällt auf die Transpiration landwirtschaftlicher und forstwirtschaftlicher Kulturen. Anpassungsstrategien an die zunehmende Trockenheit zielen darauf ab, die Ausnutzung der knappen Wasserressourcen durch Landnutzungswechsel und angepasste land- und forstwirtschaftliche Bewirtschaftung zu optimieren. Dieser Artikel gibt einen Überblick über das Potential dieser Optionen angesichts der aktuellen und der prognostizierten klimatischen Bedingungen in Nordostdeutschland.

de l'Allemagne. La recharge annuelle de la nappe est presque zéro dans quelques parts de la région. En outre, les nappes phréatiques et les surfaces de l'eau des lacs se sont baissées pendant les dernières trois décennies à beaucoup d'endroits. On suppose que ce trend va se continuer pendant les prochaines décennies. La transpiration des arbres et des plantes agricoles est la plus grande consommation de l'eau. Ainsi la production forestière et agricole doit s'adapter par optimisation de l'exploitation et par des changements de l'utilisation du sol. Cet article résume des résultats des investigations à ce sujet sous des conditions climatique actuelles et futures dans la région.

*Prof. Dr. Gunnar Lischeid, Dipl.-Hydrologe  
Marco Natkhanin, Institute of Landscape Hydrology,  
Leibniz Centre for Agricultural Landscape Research,  
Eberswalder Str. 84, 15374 Müncheberg, Germany,  
lischeid@zalf.de, marcom@zalf.de*

*Résumé: Les options du changement de l'utilisation du sol pour atténuer la pénurie en eau dans le nord-est d'Allemagne – une vue d'ensemble*

De plus en plus, le manque d'eau est considéré comme un problème grave dans les régions du nord-est

Manuscript submitted: 01/02/2010  
Accepted for publication: 10/03/2011